RESPIRONICS

Disinfection Guide for Professional Use Only

The following disinfection methods are validated for Respironics multi-patient use masks in the hospital/institutional environment.

High-Level Disinfection

Product	Thermal					Chemical						Gas Plasma	Autoclave	Max Cycles	Notes
	70° C for 100 minutes	75° C for 30 minutes	80° C for 10 minutes	90° C for 1 minute	93° C for 10 minutes	Cidex	Cidex OPA	Wavicide	Control III	Anioxyde 1000 †	Gigasept FF †	Sterrad 100S	121°C for 15 minutes		
PerformaTrak		6		6		6	6	6	6			;		10	Remove the forehead pad from the frame before disinfection. Replace the forehead pad for each patient use.
PerforMax	\oslash	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\bigotimes	\oslash	Ş	10/5	Mask may be autoclaved using gravity displacement cycle, no sterilization bag (5 cycles max). Mask may be disinfected using all other methods (10 cycles max). Anioxyde 1000 † and Gigasept FF † are approved treatments for small and large sizes only.
Total	6					\bigotimes	\oslash					\oslash		10	Approved at 70°C for 30 minutes only.

Pre-Treatment Cleaning

A Caution

Follow all instructions from the manufacturer of the pre-treatment products. Any deviation from these instructions may impact the performance of the product. Review all applicable instructions for additional warnings and cautions.

- Dissasemble the mask according to the Instructions for Use included with the product.
- Clean the mask using a soft bristle brush to adequately remove adhering substances from each component while soaking in a commercially available, anionic detergent (Examples: MEDIZIME LF, ANIOSYME DD1+). Extra attention should be given to the crevices and cavities during cleaning.
- If compatible with the manufacturers instructions, the detergent may be applied within the washer/disinfection cycle.
- Rinse the mask with 5 liters of water and air dry out of direct sunlight.

2 High-Level Disinfection

The recommended disinfection methods are identified by mask product (above chart) and approved (\oslash) for use on the mask and/or mask parts. Review the notes (=) section for exceptions or deviations.

Caution Follow all instructions from the manufacturer of the disinfection products. Any deviation from the manufacturer's instructions or use of agents other than a cautions. those listed in this guide may impact the performance or durability of the product. Review all applicable instructions for additional warnings and cautions.

/ Warnings

Before Disinfection

- The fabric materials (eg: headgear, straps) cannot be disinfected using the methods listed. The fabric materials must be replaced before multi-patient use.
- Masks with port caps: Open or remove the port cap prior to disinfecting the mask.

After Disinfection

- Inspect all parts for damage or wear; replace any parts that have visibly deteriorated (cracking, crazing, tears, etc.)
- Rinse thoroughly with water and air dry out of direct sunlight. Make sure the mask is dry before use.
- Verify the entrainment valve functions correctly, as outlined in the Instructions for Use included with the product.

Note

PHILIPS

- After disinfection, discoloration and a slight odor is normal.
- † Anioxyde 1000, Aniosyme DD1, and Gigasept FF are not available for use in the United States.

Chemical Agent Active Ingredients

Cidex - 2.4% glutaraldehyde Cidex OPA - 0.55% ortho-phthaladehyde Wavicide - 2.65% glutaraldehyde Control III - 10% alkyl dimethyl benzyl ammonium chloride, 10% alkyl dimethyl ethyl benzyl ammonium chloride Anioxyde 1000 - 0.15% peracetic acid Gigasept FF - 11.0 g succindialdehyde, 3.0 g dimethoxytetrahydrofuran

Contact Philips Respironics

Visit us on-line at www.philips.com/NIVmasks to download copies of the Disinfection Guide or contact Philips Respironics Customer Service. Respironics: 1-800-345-6443 (USA or Canada only) or 724-387-4000 Respironics Deutschland: +49 (0) 8152 93060

PerformaTrak, PerforMax, and Total are trademarks of Philips Respironics Inc.

CIDEX and STERRAD are trademarks of Johnson & Johnson. WAVICIDE and MEDIZIME LF are trademarks of Medical Chemical Corporation. CONTROL III is a trademark of Maril Products, Inc. ANIOXYDE and ANIOSYME DD1 are trademarks of Anios Laboratories, Gigasept FF is a registered trademark of Schülke&Mayr.



Respironics Inc. 1001 Murry Ridge Lane Murrysville, PA 15668 USA Respironics Deutschland Gewerbestrasse 17 82211 Herrsching, Germany ECREP



